(To be updated at least annually and as needed)



For children in childcare, kindergarten, family day care and out of school hours care

Instructions

- To be completed by parents/guardians in consultation with their child's doctor.
- Parents/guardians should inform their child's childcare service, kindergarten, family day care or out of school hours care immediately if there are any changes to this record.
- Please tick the appropriate boxes or print your responses in the blank spaces where indicated

	Method (e.g. puffer & spacer)		nd how much? puff morning and night)	
EQUIREMENTS (JSUALLY TAKEN IN CARE		_	
Does the child take any asthma medication before exercise/play?				
		☐ Other (Please specify)	
Other (Please specify) Other (Please specify)		Pets		
•	☐ Difficulty speaking	Smoke	☐ Smoke	
-	☐ Difficulty breathing	☐ Dust	☐ Dust	
	☐ Coughing	_	☐ Pollens	
chest	☐ Tightness in chest	☐ Colds/\	☐ Colds/Viruses	
	☐ Wheeze	☐ Exercis	е	
	Increased signs of:			
	Signs of child's asthma worseni	ng What trigge	rs the child's asthma	
nber: ∐ Yes □	No Membership number:			
	•			
•	,			
act telephone: (H)(W)		РНОТО	
,	•			
	(W) (M)			
an's Name:		ſ		
AILS				
(First Name)	(Family Name)			
	Sex:		of birth://	
	an's Name:	an's Name:	AILS an's Name: (W) (M) (M) (M) (M) (M) (M) (M) (M) (M) (M	

Name of Medication (e.g. Ventolin, Flixotide)	Method (e.g. puffer & spacer)	When and how much? (e.g. one puff morning and night)

ASTHMA FIRST AID PLAN

Please tick your preferred Asthma First Aid Plan ☐ 4 STEP ASTHMA FIRST AID PLAN Step 1. Sit the person upright, be calm and reassuring. Do not leave them alone. Give 4 separate puffs of a blue reliever * Step 2. The medication is best given one puff at a time via a spacer device. If a child is under 5 years of age a face mask should be used Ask the person to take 4 breaths from the spacer after each puff of medication. If a spacer and face mask is not available, use the blue reliever puffer on its own. Step 3. Wait 4 minutes. Step 4 If there is little or no improvement repeat steps 2 and 3. If there is still little or no improvement call an ambulance immediately (DIAL 000). Continue to repeat steps 2 and 3 while waiting for the ambulance. *A Bricanyl Turbuhaler may be used in first aid treatment if a puffer and spacer are unavailable. OR □ CHILD'S ASTHMA FIRST AID PLAN (approved by doctor) (if different from above) If the child's condition suddenly deteriorates or if at any time you are concerned — call an ambulance immediately (000). In the event of an asthma attack, I agree to my child receiving the treatment described above. I authorise children's services staff to assist my child with taking asthma medication should he/she require help. I will notify you in writing if there are any changes to these instructions. I agree to pay all expenses incurred for any medical treatment deemed necessary. Please notify me if my child has received asthma first aid. Parent's/Guardian's Signature: Date___/__/

For further information please contact The Asthma Foundation of Victoria on (03) 9326 7088, toll free 1800 645 130, or visit our website www.asthma.org.au

Doctor's Signature: